

HOMEWORK

- Section 4.3 - 1, 4, 6, 7, 9, 10, 11, 14, 16, 41, 42, 44, 45, 48, 52, 54, 57, 58, 59, 73, 76

SECTION 4.3 - GAUSS-JORDAN ELIMINATION

Reduced Matrices. In the last section, our goal was to reduce matrices to one of the following forms

$$\left[\begin{array}{cc|c} 1 & 0 & m \\ 0 & 1 & n \end{array} \right] \quad \left[\begin{array}{cc|c} 1 & m & n \\ 0 & 0 & 0 \end{array} \right] \quad \left[\begin{array}{cc|c} 1 & m & n \\ 0 & 0 & p \end{array} \right]$$

where m, n, p are real numbers and $p \neq 0$.

These are all examples of *reduced matrices*, or *reduced row echelon form* matrices. If the matrix has a larger size, we can still put it in reduced form, but it is hard to list out all the possibilities, so we will give a definition here

Definition 1 (Reduced Form). *A matrix is in reduced form if*

- (1) *Each row consisting entirely of zeros is below any row having at least one nonzero element.*
- (2) *The leftmost nonzero element in each row is 1.*
- (3) *All other elements in the column containing the leftmost 1 of a given row are zeros.*
- (4) *The leftmost 1 in any row is to the right of the leftmost 1 in the row above.*

Here are a few examples of matrices in reduced form

$$\left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & -3 \end{array} \right] \quad \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 9 \\ 0 & 0 & 1 & 4 \end{array} \right] \quad \left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{array} \right]$$

$$\left[\begin{array}{cccc|c} 1 & 4 & 0 & 0 & 7 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 4 \end{array} \right] \quad \left[\begin{array}{ccc|c} 1 & 0 & 4 & 8 \\ 0 & 1 & 1 & 9 \\ 0 & 0 & 0 & 2 \end{array} \right]$$

Example 1. *Why are the following matrices not in reduced form? Put them in reduced form:*

(a)

$$\left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 3 & -6 \end{array} \right]$$

(b)

$$\left[\begin{array}{ccc|c} 1 & 5 & 4 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

(c)

$$\left[\begin{array}{ccc|c} 0 & 1 & 0 & -3 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 \end{array} \right]$$

(d)

$$\left[\begin{array}{ccc|c} 1 & 2 & 0 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 4 \end{array} \right]$$

Solution.

- (a) Here, we do not have a 1 in the bottom left, we have a 3. Just divide the second row by 3, $\frac{1}{3}R_2 \rightarrow R_2$

$$\left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & -2 \end{array} \right]$$

- (b) In this one, there is not zeros above the 1 in the second row, second column. To fix this, we use $R_1 - 5R_2 \rightarrow R_1$

$$\left[\begin{array}{ccc|c} 1 & 0 & -2 & -2 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

- (c) Here, the 1's are in the wrong place. To fix it, use $R_1 \leftrightarrow R_2$

$$\left[\begin{array}{ccc|c} 0 & 1 & 0 & -3 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 \end{array} \right]$$

- (d) Here, the row with all zeros isn't below all the other rows, so to fix it, we use $R_2 \leftrightarrow R_3$

$$\left[\begin{array}{ccc|c} 1 & 2 & 0 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 4 \end{array} \right]$$

Let's now actually use Gauss-Jordan elimination to solve a system

Example 2. Solve the following system using Gauss-Jordan elimination:

$$\begin{array}{rrcr} 3x & + & y & - & 2z & = & 2 \\ x & - & 2y & + & z & = & 3 \\ 2x & - & y & - & 3z & = & 3 \end{array}$$

Solution. First we turn it into an augmented matrix

$$\left[\begin{array}{ccc|c} 3 & 1 & -2 & 2 \\ 1 & -2 & 1 & 3 \\ 2 & -1 & -3 & 3 \end{array} \right]$$

Now we follow the process to get a reduced form. Start by getting a 1 in the top left:

$$\left[\begin{array}{ccc|c} 3 & 1 & -2 & 2 \\ 1 & -2 & 1 & 3 \\ 2 & -1 & -3 & 3 \end{array} \right] \xrightarrow[R_1 \leftrightarrow R_2]{} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 3 \\ 3 & 1 & -2 & 2 \\ 2 & -1 & -3 & 3 \end{array} \right]$$

and now get zeros everywhere else in that column

$$\left[\begin{array}{ccc|c} 1 & -2 & 1 & 3 \\ 3 & 1 & -2 & 2 \\ 2 & -1 & -3 & 3 \end{array} \right] \xrightarrow[R_2 - 3R_1 \rightarrow R_2]{} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 3 \\ 0 & 7 & -5 & -7 \\ 2 & -1 & -3 & 3 \end{array} \right] \xrightarrow[R_3 - 2R_1 \rightarrow R_3]{} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 3 \\ 0 & 7 & -5 & -7 \\ 0 & 3 & -5 & -3 \end{array} \right]$$

Now we need to get a 1 in the second column, second row

$$\left[\begin{array}{ccc|c} 1 & -2 & 1 & 3 \\ 0 & 7 & -5 & -7 \\ 0 & 3 & -5 & -3 \end{array} \right] \xrightarrow[R_2 - 2R_3 \rightarrow R_2]{} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 3 \\ 0 & 1 & 5 & -1 \\ 0 & 3 & -5 & -3 \end{array} \right]$$

And now we will get 0's in the other entries in the second column

$$\left[\begin{array}{ccc|c} 1 & -2 & 1 & 3 \\ 0 & 1 & 5 & -1 \\ 0 & 3 & -5 & -3 \end{array} \right] \xrightarrow[R_1 + 2R_2 \rightarrow R_1]{} \left[\begin{array}{ccc|c} 1 & 0 & 11 & 1 \\ 0 & 1 & 5 & -1 \\ 0 & 3 & -5 & -3 \end{array} \right] \xrightarrow[R_3 - 3R_2 \rightarrow R_3]{} \left[\begin{array}{ccc|c} 1 & 0 & 11 & 1 \\ 0 & 1 & 5 & -1 \\ 0 & 0 & -20 & 0 \end{array} \right]$$

Now, we get a 1 in the bottom left

$$\left[\begin{array}{ccc|c} 1 & 0 & 11 & 1 \\ 0 & 1 & 5 & -1 \\ 0 & 0 & -20 & 0 \end{array} \right] \xrightarrow[-\frac{1}{20}R_3 \rightarrow R_3]{} \left[\begin{array}{ccc|c} 1 & 0 & 11 & 1 \\ 0 & 1 & 5 & -1 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

And finally we get 0's everywhere else in that column

$$\left[\begin{array}{ccc|c} 1 & 0 & 11 & 1 \\ 0 & 1 & 5 & -1 \\ 0 & 0 & 1 & 0 \end{array} \right] \xrightarrow[R_1 - 11R_3 \rightarrow R_1]{} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 5 & -1 \\ 0 & 0 & 1 & 0 \end{array} \right] \xrightarrow[R_2 - 5R_3 \rightarrow R_2]{} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

So, the solution is $x = 1$, $y = -1$, and $z = 0$.

Example 3. Solve by Gauss-Jordan elimination:

$$\begin{array}{rrcr} 2x_1 & - & 4x_2 & - & x_3 & = & -8 \\ 4x_1 & - & 8x_2 & + & 3x_3 & = & 4 \\ -2x_1 & + & 4x_2 & + & x_3 & = & 11 \end{array}$$

Solution. Begin by finding the augmented matrix

$$\left[\begin{array}{ccc|c} 2 & -4 & -1 & -8 \\ 4 & -8 & 3 & 4 \\ -2 & 4 & 1 & 11 \end{array} \right]$$

Now we get the 1 in the upper left

$$\left[\begin{array}{ccc|c} 2 & -4 & -1 & -8 \\ 4 & -8 & 3 & 4 \\ -2 & 4 & 1 & 11 \end{array} \right] \xrightarrow[\sim]{\frac{1}{2}R_1 \rightarrow R_1} \left[\begin{array}{ccc|c} 1 & -2 & -\frac{1}{2} & -4 \\ 4 & -8 & 3 & 4 \\ -2 & 4 & 1 & 11 \end{array} \right]$$

Now we get the zeros in the rest of the column

$$\left[\begin{array}{ccc|c} 1 & -2 & -\frac{1}{2} & -4 \\ 4 & -8 & 3 & 4 \\ -2 & 4 & 1 & 11 \end{array} \right] \xrightarrow[\sim]{R_2 - 4R_1 \rightarrow R_2} \left[\begin{array}{ccc|c} 1 & -2 & -\frac{1}{2} & -4 \\ 0 & 0 & 5 & 20 \\ -2 & 4 & 1 & 11 \end{array} \right] \xrightarrow[\sim]{R_3 + 2R_1 \rightarrow R_3} \left[\begin{array}{ccc|c} 1 & -2 & -\frac{1}{2} & -4 \\ 0 & 0 & 5 & 20 \\ 0 & 0 & 0 & 3 \end{array} \right]$$

The last row of the matrix now corresponds to an equation of the form $0 = 3$, which is nonsense. Thus this system has no solution.

Example 4. Solve by Gauss-Jordan elimination:

$$\begin{array}{rrcr} 3x_1 & + & 5x_2 & - & x_3 & = & -7 \\ x_1 & + & x_2 & + & x_3 & = & -1 \\ 2x_1 & & & + & 11x_3 & = & 7 \end{array}$$

Solution. $x_1 = -2, x_2 = 0, x_3 = 1$

Example 5. Solve by Gauss-Jordan elimination:

$$\begin{array}{rrcr} 3x_1 & - & 4x_2 & - & x_3 & = & 1 \\ 2x_1 & - & 3x_2 & + & x_3 & = & 1 \\ x_1 & - & 2x_2 & + & 3x_3 & = & 2 \end{array}$$

Solution. No solution.