

INTRO TO DERIVATIVES

Math 130 - Essentials of Calculus

19 February 2021

INSTANTANEOUS RATE OF CHANGE

Before, we were trying to estimate the speed of a baseball 1 second after being thrown straight upward, where the height of the ball after t seconds was given by:

$$h(t) = 36t - 16t^2.$$

INSTANTANEOUS RATE OF CHANGE

Before, we were trying to estimate the speed of a baseball 1 second after being thrown straight upward, where the height of the ball after t seconds was given by: $h(t) = 36t - 16t^2$. The method we used was to shrink the interval of time that we took the average over.

INSTANTANEOUS RATE OF CHANGE

Before, we were trying to estimate the speed of a baseball 1 second after being thrown straight upward, where the height of the ball after t seconds was given by: $h(t) = 36t - 16t^2$. The method we used was to shrink the interval of time that we took the average over. That is, we used the process

$$\lim_{\Delta t \rightarrow 0} \frac{\Delta h}{\Delta t} = \lim_{t \rightarrow 1} \frac{h(t) - h(1)}{t - 1}.$$

INSTANTANEOUS RATE OF CHANGE

Generally speaking, the instantaneous rate of change is the limit of the average rate of change over successively smaller and smaller intervals.

INSTANTANEOUS RATE OF CHANGE

Generally speaking, the instantaneous rate of change is the limit of the average rate of change over successively smaller and smaller intervals.

DEFINITION (INSTANTANEOUS RATE OF CHANGE)

The instantaneous rate of change of a function f at the input value x_1 is

$$\lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \lim_{x_2 \rightarrow x_1} \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

provided the limit exists.

INSTANTANEOUS RATE OF CHANGE

An alternative, but equivalent definition for the instantaneous rate of change is

DEFINITION (INSTANTANEOUS RATE OF CHANGE)

The instantaneous rate of change of a function f at the input value a is

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

provided the limit exists.

Think of $x_1 = a$, $x_2 = a + h$, then $\Delta x = h$.


INSTANTANEOUS RATE OF CHANGE

EXAMPLE

A rock is dropped from a bridge over a river. The distance, in meters, between the rock and the river t seconds after the rock is dropped is given by $s(t) = 48 - 4.9t^2$. Compute the speed of the rock after 2 seconds.


SLOPE OF THE TANGENT LINE

Recall that the average rate of change was the slope of a *secant line*. As we shrink the interval that the average rate of change is taken over, the slope of the secant lines approaches the slope of the tangent line.




SLOPE OF THE TANGENT LINE

Recall that the average rate of change was the slope of a *secant line*. As we shrink the interval that the average rate of change is taken over, the slope of the secant lines approaches the slope of the tangent line.




SLOPE OF THE TANGENT LINE

Recall that the average rate of change was the slope of a *secant line*. As we shrink the interval that the average rate of change is taken over, the slope of the secant lines approaches the slope of the tangent line.




SLOPE OF THE TANGENT LINE

Recall that the average rate of change was the slope of a *secant line*. As we shrink the interval that the average rate of change is taken over, the slope of the secant lines approaches the slope of the tangent line.



SLOPE OF THE TANGENT LINE

Recall that the average rate of change was the slope of a *secant line*. As we shrink the interval that the average rate of change is taken over, the slope of the secant lines approaches the slope of the tangent line.



SLOPE OF THE TANGENT LINE

DEFINITION (SLOPE OF TANGENT LINE)

The tangent line to the curve $y = f(x)$ at the point $(x_1, f(x_1))$ is the line through this point with slope

$$m = \lim_{x_2 \rightarrow x_1} \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

provided the limit exists.

SLOPE OF THE TANGENT LINE

DEFINITION (SLOPE OF TANGENT LINE)

The tangent line to the curve $y = f(x)$ at the point $(x_1, f(x_1))$ is the line through this point with slope

$$m = \lim_{x_2 \rightarrow x_1} \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

provided the limit exists.

DEFINITION (SLOPE OF TANGENT LINE)

The tangent line to the curve $y = f(x)$ at the point $(a, f(a))$ is the line through this point with slope

$$m = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

provided the limit exists.

FINDING THE TANGENT LINE

EXAMPLE

Find the equation of the tangent line to the given function at the given point:

❶ $y = 2x^2 + 1$ at $(3, 19)$

FINDING THE TANGENT LINE

EXAMPLE

Find the equation of the tangent line to the given function at the given point:

❶ $y = 2x^2 + 1$ at $(3, 19)$

❷ $f(x) = 3x - x^2$ at $(1, 2)$

THE DERIVATIVE

The slope, or instantaneous rate of change, of a function is usually referred to as *the derivative* of the function, denoted $f'(a)$.

THE DERIVATIVE

The slope, or instantaneous rate of change, of a function is usually referred to as *the derivative* of the function, denoted $f'(a)$.

DEFINITION (DERIVATIVE)

The derivative of the function f at the number a , denoted $f'(a)$, is defined by

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

if the limit exists.

Likewise, we could use the alternate form of the difference quotient to compute a derivative as well

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}.$$

THE DERIVATIVE

EXAMPLE

Compute the derivatives of the following functions at the given value:

❶ $g(t) = t^2 + 4t, t = 1$

THE DERIVATIVE

EXAMPLE

Compute the derivatives of the following functions at the given value:

❶ $g(t) = t^2 + 4t, t = 1$

❷ $f(x) = \sqrt{2x}, x = 2$